arXiv:1108.1539v1 [math.RT] 7 Aug 2011

ON THE DIMENSION OF LOCALLY ANALYTIC REPRESENTATIONS OF SEMISIMPLE p-ADIC GROUPS

TOBIAS SCHMIDT

PRELIMINARY VERSION

Abstract. We prove an analogue of a theorem of S.P. Smith on universal enveloping algebras ([21]) in the context of locally analytic representation theory. Let G be a p-adic Lie group whose Lie algebra is split semisimple. Let r be half the dimension of a minimal co-adjoint orbit. The values of r are well-known. We show that the canonical dimension of any admissible locally analytic G-representation which is not zero-dimensional is at least r. We prove this result by reducing it to an analogous statement for modules over p-adically completed deformations of universal enveloping algebras which was established in a recent preprint by K. Ardakov and S. Wadsley ([3]).

Contents

1. Introduction 1
2. Distribution algebras of L-uniform groups 3
3. Dimensions 7
References 12

1. Introduction

Let p be an odd prime number and $\mathbb{Q}_p \subseteq L \subseteq K \subseteq \mathbb{C}_p$ a chain of finite extensions of $\mathbb{Q}_p$. Let G be a d-dimensional locally L-analytic group (e.g. the group of rational points of a linear algebraic group over L).

In a series of papers (cf. [18] for a survey) P. Schneider and J. Teitelbaum have developed a theory of admissible locally analytic G-representations. On the one hand, this theory has found substantial arithmetic applications in the field of p-adic automorphic forms, p-adic interpolation or the p-adic Langlands programme. On the other hand, it is of independent representation theoretic interest not only because it provides a unified framework for studying finite-dimensional algebraic representations as well as the admissible-smooth representations of Langlands theory. Apart from the latter two classes there are many representations which are *genuinely locally analytic*. A possible way to organize this situation and, more specifically, to meisure the 'size' of the vector space underlying a representation (which usually is of infinite vector space dimension) one may introduce an Auslander-Gorenstein style dimension function on the abelian category $Rep^a_G(K)$ of admissible locally analytic G-representations over K (cf. [19][1]). It associates a well-defined number $0 \leq \dim(V) \leq d$ to each nonzero admissible representation V. In this way one obtains a filtration of $Rep^a_G(K)$ by Serre subcategories

$$Rep^a_G(K) = Rep^a_G(K)_d \supseteq Rep^a_G(K)_{d-1} \supseteq ... \supseteq Rep^a_G(K)_1 \supseteq Rep^a_G(K)_0$$

where V lies in $Rep^a_G(K)_i$ if and only if $\dim(V) \leq i$. The aforementioned algebraic representations and the smooth-admissible representations are concentrated in dimension zero.

In this note, we investigate this situation under the additional assumption that the Lie algebra $Lie(G)$ of G is split semisimple over L. It turns out that, in this case, there is a large 'gap' in the above filtration. Namely, let $\mathbf{G}_{\mathbb{C}}$ be a complex semisimple algebraic group having the same root system as $Lie(G)$ and let r be half the smallest possible dimension of a non-zero co-adjoint $\mathbf{G}_{\mathbb{C}}$-orbit. The values of r are well-known (cf. 3.5 below). Our main result says that if V is an admissible locally analytic G-representation over K which is not zero-dimensional then $\dim(V) \geq r$. Moreover, we show that an admissible locally analytic G-representations is zero-dimensional if and only if its associated coherent sheaf (in the sense of [19]) consists of finite-dimensional K-vector spaces.

We remark that the above theorem was inspired by a classical theorem of S.P. Smith on universal enveloping algebras of semisimple Lie algebras ([21]) as well as a p-adically completed version of this theorem established in a recent preprint by K. Ardakov and S. Wadsley ([3]). In fact, the results on locally analytic distribution algebras as obtained in [15] and [16] enable us to make a rather straightforward reduction to the case treated in [3].

The author is grateful to K. Ardakov for some useful discussions concerning the work [3].

Notation: Let p be an odd prime. Let $\mathbb{Q}_p \subseteq L \subseteq K \subseteq \mathbb{C}_p$ be a chain of finite extensions of $\mathbb{Q}_p$. The absolute value $|.|$ on $\mathbb{C}_p$ is normalized as usual by $|p| = p^{-1}$. Let $o_L \subseteq L$ be the ring of integers and $\pi_L \in o_L$ a uniformizer. Let n and e be the degree and the ramification index of the extension $L/\mathbb{Q}_p$ respectively. Also, $o_K \subset K$ denotes the integers in K and $\pi_K \in o_K$ a uniformizer. Let $k := o_K/(\pi_K)$ denote the residue field of K.
An o_K-submodule of a vector space V over K is called a *lattice* if it contains a basis of V over K.

---

[1] In [loc.cit.] the authors consider the associated *codimension* function.

2. DISTRIBUTION ALGEBRAS OF L-UNIFORM GROUPS

2.1. We refer to [6] for the definition and a thorough discussion of uniform pro-p groups. Let G be a uniform pro-p group of dimension $d := \dim_{\mathbb{Q}_p} G$. Let $h_1, ..., h_d$ be a set of topological generators. In particular, any element $g \in G$ can uniquely be written as $g = h_1^{\lambda_1} \cdots h_d^{\lambda_d}$ with $\lambda_1, ..., \lambda_d \in \mathbb{Z}_p$. According to [loc.cit.], Thm. 4.30 the operations

$$\begin{aligned} \lambda \cdot x &= x^{\lambda}, \\ x + y &= \lim_{i\to\infty}(x^{p^i} y^{p^i})^{p^{-i}}, \\ [x, y] &= \lim_{i\to\infty}(x^{-p^i} y^{-p^i} x^{p^i} y^{p^i})^{p^{-2i}} \end{aligned}$$

for $x, y \in G$ and $\lambda \in \mathbb{Z}_p$ define on the set G the structure of a Lie algebra L_G over $\mathbb{Z}_p$. This Lie algebra is *powerful* in the sense that it is a free $\mathbb{Z}_p$-module (of finite rank d) and satisfies $[L_G, L_G] \subseteq pL_G$ (recall $p \neq 2$).

Let $Lie(G)$ denote the Lie algebra of the locally $\mathbb{Q}_p$-analytic manifold G and

$$exp_G : Lie(G) - - > G$$

its exponential map. There is a canonical inclusion of $\mathbb{Z}_p$-Lie algebras $L_G \subset Lie(G)$ such that $\exp_G$ is defined on L_G and maps it bijectively onto G. Moreover, the $\mathbb{Z}_p$-lattice $L_G \subset Lie(G)$ is uniquely determined by these two properties ([loc.cit.], sect. 4.5/9.4).

2.2. Now let G be a compact locally L-analytic group of dimension $d = \dim_L G$. We will occasionally write G_0 for its locally $\mathbb{Q}_p$-analytic structure induced by scalar restriction to $\mathbb{Q}_p$ whenever we need to emphasize this. We point out that there is a canonical isomorphism of $\mathbb{Q}_p$-Lie algebras $Lie(G) \simeq Lie(G_0)$ ([4], 5.14.5) which we will use to identify both $\mathbb{Q}_p$-algebras from now on. Following [15],[13] we call G *L-uniform* if

1. G is uniform pro-p,
2. the $\mathbb{Z}_p$-submodule $L_G \subset Lie(G)$ is stable under multiplication by o_L (and so is an o_L-submodule of $Lie(G)$).

We remark that any compact locally L-analytic group G has a basis of neighbourhoods of $1 \in G$ consisting of open normal subgroups which are L-uniform ([15], Cor.4.4).

Let in the following G be a locally L-analytic group of dimension d which is L-uniform. Then L_G is a free o_L-module of rank d. Let $\mathfrak{x}_1, ...\mathfrak{x}_d$ be an o_L-basis of L_G and let $v_1, ..., v_n$ be a $\mathbb{Z}_p$-basis of o_L with the additional property $v_1 = 1$. Then the elements

$$h_{ij} := \exp_G(v_i \mathfrak{x}_j)$$

are a set of topological generators of G of cardinality $nd = \dim_{\mathbb{Q}_p} G_0$.

2.3. For the definition and main properties of locally analytic distribution algebras we refer to [19]. Let $D(G) := D(G, K)$ be the algebra of K-valued locally L-analytic distributions on G and define $D(G_0)$ similarly for G_0. These are topological K-Fréchet algebras and the inclusion of locally L-analytic functions into locally $\mathbb{Q}_p$-analytic functions on G gives rise to a quotient map of topological algebras

$$D(G_0) \longrightarrow D(G)$$

whose kernel ideal is generated by the kernel of the natural map

$$L \otimes_{\mathbb{Q}_p} Lie(G_0) \longrightarrow Lie(G), a \otimes \mathfrak{x} \mapsto a\mathfrak{x} \tag{2.3}$$

([15], lemma 5.1).

The underlying K-vector space of $D(G_0)$ admits the following description. Fix an ordering of the generators h_{ij}. Let $b_{ij} := h_{ij} - 1 \in \mathbb{Z}[G]$ and $\mathbf{b}^\alpha := \prod_{ij} b_{ij}^{\alpha_{ij}} \in \mathbb{Z}[G]$ (in the given ordering) for $\alpha \in \mathbb{N}_0^{nd}$. Then $D(G_0)$ equals the set of all convergent series

$$\lambda = \sum_{\alpha \in \mathbb{N}_0^{nd}} d_\alpha \mathbf{b}^\alpha$$

with $d_\alpha \in K$ such that the set $\{|d_\alpha| r^{|\alpha|}\}_\alpha$ is bounded for all real numbers $0 < r < 1$. The family of norms $||.||_r,\ 0 < r < 1$ where

$$||\lambda||_r := \sup_\alpha |d_\alpha| r^{|\alpha|}$$

respectively their quotient norms define the Fréchet topologies on $D(G_0)$ respectively $D(G)$. The norms belonging to the interval $p^{-1} \leq r < 1$ where $r \in p^{\mathbb{Q}}$ are multiplicative. In this case the corresponding norm completions $D_r(G_0)$ and $D_r(G)$ of $D(G_0)$ and $D(G)$ respectively are K-Banach algebras. Furthermore, we emphasize that the norms in this case do not depend on the choice of minimal set of topological generators for the group G ([19], discussion after Thm. 4.10).

2.4. Let $r \in p^{\mathbb{Q}}$ such that $p^{-1} \leq r < 1$. There is the following norm filtration on $D_r(G_0)$ defined by the additive subgroups

$$\begin{aligned} F_r^s D_r(G_0) :=& \ \ \{\lambda \in D_r(G_0),\ ||\lambda||_r \leq p^{-s}\}, \\ F_r^{s+} D_r(G_0) :=& \ \ \{\lambda \in D_r(G_0),\ ||\lambda||_r < p^{-s}\} \end{aligned}$$

for $s \in \mathbb{R}$ with graded ring

$$gr_r^\bullet D_r(G_0) := \oplus_{s \in \mathbb{R}}\ gr_r^s D_r(G_0)$$

where $gr_r^s D_r(G_0) := F_r^s D_r(G_0) / F_r^{s+} D_r(G_0)$.

We may use the quotient norm of $||.||_r$ to define a norm filtration on $D_r(G)$ as above. It coincides with the quotient filtration of $F_r^\bullet D_r(G_0)$. We denote the associated graded ring by $gr_r^\bullet D_r(G)$.
Similarly, the absolute value on K induces a filtration and a graded ring grK as above. The choice of uniformizer π_K induces an isomorphism of grK with the ring of Laurent polynomials in one variable over k. The rings $gr_r^\bullet D_r(G_0)$ and $gr_r^\bullet D_r(G)$ are naturally grK-algebras.

Now let additionally $p^{-1} < r < 1$. Then Thm. 4.5 of [19] shows that each $gr^{\bullet}_r D_r(G_0)$ equals a polynomial ring over grK in nd variables (the principal symbols of the b_{ij}). Hence any $gr^{\bullet}_r D_r(G)$ will be a commutative grK-algebra.

2.5. Let $\{G^{p^m}\}_{m \geq 0}$ be the lower p-series of the uniform pro-p group G, i.e.

$$G^{p^m} := \{g^{p^m} : g \in G\}$$

for $m \geq 0$. Each G^{p^m} is a characteristic open uniform pro-p subgroup of G of index p^{mnd}. The set $\{G^{p^m}\}_m$ constitutes a neighbourhood basis of $1 \in G$. Because of $L_{G_0^{p^m}} = p^m L_{G_0}$ each open subgroup G^{p^m}, endowed with its induced locally L-analytic structure, is L-uniform. By functoriality of $D(\cdot)$ ([8]) we obtain a series of subalgebras

$$D(G_0) \supseteq D(G_0^p) \supset D(G_0^{p^2}) \supset ...$$

which yields, by passage to quotients, a series

$$D(G) \supseteq D(G^p) \supset D(G^{p^2}) \supset ...$$

of subalgebras of $D(G)$. For each $m \geq 0$ we may apply the above discussion to the group G^{p^m} and obtain a family of norms $||.||_r^{(m)}$ on $D(G_0^{p^m})$ as well as quotient norms q_r^m on $D(G^{p^m})$. Of course, these norms again induce filtrations and graded rings as explained above whenever $r \in p^{\mathbb{Q}}$ with $p^{-1} \leq r < 1$.

To ease notation we put for $m \geq 0$ from now on

$$r_m := \sqrt[p^m]{1/p}.$$

In particular, $r_0 = 1/p$.

Lemma 2.6. *The ring extension $D(G^{p^m}) \subset D(G)$ completes to a ring extension*

$$D_{r_0}(G^{p^m}) \subset D_{r_m}(G)$$

and $D_{r_m}(G)$ is a finite and free (left or right) module over $D_{r_0}(G^{p^m})$ on a basis any system of coset representatives for the group G/G^{p^m}.

Proof. This is a slight generalization of [15], Lemma 7.4 using the fact that the norm $||.||_{r_m}^{(0)}$ on $D(G_0)$ restricts to $||.||_{r_0}^{(m)}$ on the subring $D(G_0^{p^m}) \subset D(G_0)$. This latter fact is itself a mild extension of [loc.cit.], Prop. 6.2 which follows with the same proof given there. □

2.7. We remind the reader ([12], 1.5.8) that a *crossed product* of a ring S by a group H is an associative ring $S * H$ which contains S as a subring and contains a set of units $\overline{H} = \{\overline{h} : h \in H\}$, isomorphic as a set to H, such that

(i) $S * H$ is a free left S-module with basis $\overline{H}$;
(ii) for all $x, y \in \overline{H}, \overline{x}S = S\overline{x}$ and $\overline{x} \cdot \overline{y}S = \overline{xy}S$.

Given such a crossed product one obtains maps $\sigma : H \to \operatorname{Aut}(S)$ (an *action*) and $\tau : H \times H \to S^\times$ (a *twisting*) by the rules $\sigma(x)(s) = \overline{x}^{-1} s \overline{x}$ and $\overline{x} \cdot \overline{y} = \overline{xy}\tau(x, y)$. It follows that σ defines a group homomorphism $H \to \operatorname{Out}(S)$ and that τ is a 2-cocycle for the natural action of H on $S^\times$ via σ. Conversely, starting with a ring S, a group H, a group homomorphism $\sigma : H \to \operatorname{Out}(S)$ and a 2-cocycle $\tau : H \times H \to S^\times$ one can construct an associative ring $S *_{\sigma,\tau} H$ which is a crossed product of S by H having the prescribed action and twisting ([14]).

By elementary properties of $D(\cdot)$ the ring $D(G)$ is a crossed product of the ring $S := D(G^{p^m})$ by the group $H := G/G^{p^m}$ with action and twisting given as follows. Fix a set of representatives $\overline{H} \subset G \subset D(G)^\times$ for the group G/G^{p^m}. Then $\sigma(h)(s) = \delta_h^{-1} s \delta_h$ where δ_h denotes the Dirac distribution in $D(G)$ associated to $h \in \overline{H}$. On the other hand, given $h_1, h_2 \in H$ we have

$$\delta_{h_1}\delta_{h_2} = \delta_{h_1 h_2} = \delta_{h_3}\tau(h_1, h_2)$$

with some uniquely determined $h_3 \in \overline{H}, \tau(h_1, h_2) \in G^{p^m}$ and the resulting $\tau : H \times H \to G^{p^m} \subseteq S^\times$ equals the twisting.

As we have pointed out the norm $||.||_{r_0}^{(m)}$ on $S = D(G^{p^m})$ is independent of the choice of topological generators for the group G^{p^m}. This means that for $h \in G$ the conjugation automorphism on $S = D(G^{p^m})$ given by $x \mapsto \delta_h^{-1} x \delta_h$ is an isometry. Thus, σ extends to an action $H \to \operatorname{Out}(D_{r_0}(G^{p^m}))$. Likewise τ extends to a 2-cocycle with values in $D_{r_0}(G^{p^m})^\times$ and we may form the crossed product

$$D_{r_0}(G^{p^m}) *_{\sigma,\tau} H.$$

Corollary 2.8. *There is a natural K-algebra isomorphism*

$$D_{r_0}(G^{p^m}) *_{\sigma,\tau} G/G^{p^m} \xrightarrow{\cong} D_{r_m}(G).$$

Proof. The preceding lemma gives a K-algebra homomorphism $D_{r_0}(G^{p^m}) *_{\sigma,\tau} G/G^{p^m} \to D_{r_m}(G)$ induced from the inclusion $D_{r_0}(G^{p^m}) \subset D_{r_m}(G)$ which is bijective.

□

Remark: In the case $L = \mathbb{Q}_p$ this answers a question in [3], 1.5.

2.9. We fix $m \geq 1$ and consider the filtration on the Banach algebra $D_{r_0}(G^{p^m})$ induced by the norm $q_{r_0}^m$. The nd elements $h_{ij}^{p^m}$ are a minimal ordered set of topological generators for the group G^{p^m}. Define the d elements $c_j := h_{1j}^{p^m} - 1$ and let $\mathbf{c}^\beta = \prod_j c_j^{\beta_j}$ for $\beta \in \mathbb{N}^d$.

Lemma 2.10. *Let $m \geq 1$. The graded ring $gr_{r_0}^\bullet D_{r_0}(G^{p^m})$ equals a polynomial algebra over grK in the principal symbols of the d elements c_j. The K-vector space underlying $D_{r_0}(G^{p^m})$ is given by all series*

$$\lambda = \sum_{\beta \in \mathbb{N}_0^d} d_\beta \mathbf{c}^\beta$$

with $|d_\beta| {r_0}^{|\beta|} \to 0$ *for* $|\beta| \to \infty$*. Moreover, the norm* $q^m_{r_0}(\lambda)$ *may be computed as*

$$q^m_{r_0}(\lambda) = \sup_\beta q^m_{r_0}(d_\beta \mathbf{c}^\beta) = \sup_\beta |d_\beta| {r_0}^{|\beta|}.$$

Proof. We have already remarked (proof of lemma 2.6) that $||.||^{(0)}_{r_m}$ restricts to $||.||^{(m)}_{r_0}$ on the subring $D(G_0^{p^m}) \subset D(G_0)$ from which we obtain an inclusion of rings

$$gr_{r_0} D_{r_0}(G_0^{p^m}) \subset gr_{r_m} D_{r_m}(G_0)$$

by left exactness of the functor gr. Since $m \geq 1$ we have $p^{-1} < r_m$. Thus the right-hand side is a commutative ring. It follows that $gr_{r_0} D_{r_0}(G_0^{p^m})$ is a polynomial algebra over grK in the nd principal symbols of the $h_{ij}^{p^m} - 1$. Arguing as in [15], Prop. 5.6/9 the quotient $gr_{r_0} D_{r_0}(G^{p^m})$ equals a polynomial algebra in the principal symbols of the c_j and the remaining statements follow from this.

□

Remark: The result generally does not hold for $m = 0$: let $L = \mathbb{Q}_p$ and $\mathbb{Z}_p[[G]] \subseteq D_{r_0}(G)$ be the completed group ring of the profinite group G. By one of the main theorems of M. Lazard's seminal paper on p-adic analytic groups ([9], Thm. III.2.3.3) the algebra $gr_{r_0}\mathbb{Z}_p[[G]]$ equals the universal enveloping algebra of a certain Lie algebra associated to the group G which, generally, is non-abelian.

Still assuming $m \geq 1$ the preceding discussion shows that the o_K-submodule $F^0_{r_0} D_{r_0}(G^{p^m})$ of $D_{r_0}(G^{p^m})$ consists of all series

$$\lambda = \sum_{\beta \in \mathbb{N}_0^d} e_\beta (c_1/p)^{\beta_1} \cdots (c_d/p)^{\beta_d}$$

with $|e_\beta| \leq 1$ and $|e_\beta| \to 0$ for $|\beta| \to \infty$. Mapping such a series to $\sum_\beta (e_\beta \bmod \pi_K) u_1^{\beta_1} \cdots u_d^{\beta_d}$ induces an isomorphism of k-algebras between

$$gr^0_{r_0} D_{r_0}(G^{p^m}) = F^0_{r_0} D_{r_0}(G^{p^m}) / \pi_K F^0_{r_0} D_{r_0}(G^{p^m})$$

and the polynomial algebra over k in the variables $u_1, ..., u_d$. We see that the o_K-subalgebra $F^0_{r_0} D_{r_0}(G^{p^m})$ is a π_K-adically separated and complete lattice in the K-algebra $D_{r_0}(G^{p^m})$. Moreover, the π_K-adic reduction of this lattice is endowed with a positive and hence complete split $\mathbb{Z}$-filtration given by total degree with respect to the variables u_j. We conclude that $D_{r_0}(G^{p^m})$ is a *complete doubly filtered* K*-algebra* in the sense of [3], Def. 3.1 with

$$Gr(D_{r_0}(G^{p^m})) := gr(gr^0_{r_0} D_{r_0}(G^{p^m})) \simeq k[u_1, ..., u_d]. \tag{2.10}$$

3. DIMENSIONS

3.1. We recall (cf. [11], chap. III.) the notion of an Auslander regular ring. Let R be an arbitrary associative unital ring. For any (left or right) R-module N the *grade* $j_R(N)$ is defined to be either the smallest integer l such that $\mathrm{Ext}^l_R(N, R) \neq 0$ or ∞. A left and right noetherian ring R is called left and right *Auslander regular* if its left and right global dimension is finite and if every finitely generated left or right R-module N satisfies *Auslander's condition*: for any $l \geq 0$ and any R-submodule $L \subseteq \mathrm{Ext}^l_R(N, R)$ one has $j_R(L) \geq l$.

In the following the term module always means *left* module. Noetherian rings are two-sided noetherian and other ring-theoretic properties such as Auslander regular are used similarly.

Let R be an Auslander regular ring and M a finitely generated R-module. The number

$$d_R(M) := gld(R) - j_R(M)$$

is called the *canonical dimension* of M. The map $M \mapsto d_R(M)$ is a finitely partitive exact dimension function in the sense of [12], §6.8.4 and §8.3.17. For this and more details on the canonical dimension for finitely generated modules over an Auslander regular ring (or more generally, an Auslander-Gorenstein ring) we refer to [10].

A commutative noetherian ring is Auslander regular if and only if it is regular ([11], III.2.4.3). Let R be an associative ring endowed with a separated exhaustive $\mathbb{Z}$-filtration by additive subgroups such that R is complete in the filtration topology. If the associated graded ring of R is Auslander regular of (left and right) global dimension d then R is Auslander regular and has (left and right) global dimension $\leq d$ ([loc.cit.], II.2.2.1, II.3.1.4, III.2.2.5).

3.2. We return to the situation of the last section and use the same notation as there. In particular, G denotes a locally L-analytic group which is L-uniform. The rings underlying the K-Banach algebras $D_{r_0}(G^{p^m})$ and $D_{r_m}(G)$ are noetherian ([19], Thm. 4.5).

Proposition 3.3. *Let $m \geq 1$ and let A denote one of the rings $D_{r_0}(G^{p^m})$ or $D_{r_m}(G)$. Then A is Auslander regular of (left and right) global dimension equal to $d = \dim_L G$. A finitely generated A-module M is finite dimensional over K if and only if $d_A(M) = 0$.*

Proof. By Lem. 2.6 the ring $D_{r_m}(G)$ is a finite free extension of $D_{r_0}(G^{p^m})$ on a basis a system of representatives for the group G/G^{p^m}. By [19], Lem. 8.8 and [15], Cor. 7.3 it therefore suffices to prove the proposition in the case $A = D_{r_0}(G^{p^m})$. By Lem. 2.10 and the discussion above the ring A is seen to be Auslander regular of global dimension $\leq d$. Now recall that A is a complete doubly filtered K-algebra with $Gr(A) = k[u_1, ..., u_d]$. Furthermore, functoriality of $D(\cdot)$ gives a K-algebra homomorphism

$$D(G^{p^m}) \to D(\{1\}) = K$$

which extends to A and hence, A admits a nonzero module of finite K-dimension. Now [3], Prop. 9.1 completes the proof. □

For future reference we repeat the following ingredient of the preceding proof.

Corollary 3.4. *If M denotes a finitely generated $D_{r_m}(G)$-module M then*

$$d_{D_{r_m}(G)}(M) = d_{D_{r_0}(G^{p^m})}(M).$$

3.5. Let $\mathbf{G}$ be a connected split semisimple affine algebraic group scheme over o_L with Lie algebra $\mathfrak{g}$. Let us assume that the odd prime p is *very good* for $\mathbf{G}$ (in the sense of [3], 6.6). For example, if $p \geq 7$ and no irreducible component of the root system of $\mathbf{G}$ is of type A_{kp-1} for some integer $k \geq 1$ then p is a very good prime for $\mathbf{G}$. Let $\mathfrak{g}_{\mathbb{C}}$ be the complex semisimple Lie algebra with the same root system as $\mathbf{G}$ and let $G_{\mathbb{C}}$ denote the adjoint complex algebraic group associated with $\mathfrak{g}_{\mathbb{C}}$. It is known that there is a unique non-zero $\mathbf{G}_{\mathbb{C}}$-orbit in $\mathfrak{g}_{\mathbb{C}}^*$ of minimal dimension ([5], Rem. 4.3.4) and its dimension is an even integer ≥ 2. Let r denote half this dimension. The values of r are well-known, the following table is taken from [21], 1.6.

Φ	A_l	B_l	C_l	D_l	E_6	E_7	E_8	F_4	G_2
dim $\mathfrak{g}_{\mathbb{C}}$	l^2+2l	$2l^2+l$	$2l^2+l$	$2l^2-l$	78	133	248	52	14
r	l	$2l-2$	l	$2l-3$	11	17	29	8	3.

3.6. Let $\tilde{G}$ be a compact locally L-analytic group of dimension d whose Lie algebra $Lie(\tilde{G})$ is split semisimple over L. Assume the group scheme $\mathbf{G}$ has the additional property that the Lie algebra $\mathfrak{g}_L := L \otimes_{o_L} \mathfrak{g}$ is isomorphic to $Lie(\tilde{G})$. Let us fix such an isomorphism once and for all. Choose $k \geq 0$ sufficiently large such that $\exp_{\tilde{G}}$ is defined on $\Lambda := p^{k+1}\mathfrak{g}$ and such that

$$G := \exp_{\tilde{G}}(\Lambda)$$

is an open subgroup of $\tilde{G}$ which is L-uniform ([13], Lem. 2.2.4). Then

$$\frac{1}{p}L_G = \frac{1}{p}\Lambda = p^k\mathfrak{g}_0$$

as Lie algebras over $\mathbb{Z}_p$ where $\mathfrak{g}_0$ equals $\mathfrak{g}$ but viewed over $\mathbb{Z}_p$. In the following we work relative to an arbitrary but fixed number $m \geq 1$ and study the $\mathbb{Z}_p$-Lie algebra

$$\mathfrak{h} := \frac{1}{p}L_{G^{p^m}} = \frac{1}{p}p^m L_G = p^m \frac{1}{p} L_G = p^{m+k}\mathfrak{g}_0. \tag{3.7}$$

Let $U(\cdot)$ be the universal enveloping algebra of whatsoever Lie algebra we wish to consider. Let

$$\widehat{U(\mathfrak{h})}_{\mathbb{Q}_p} = (\varprojlim_i U(\mathfrak{h})/p^i U(\mathfrak{h})) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$$

be the p-adic completion of $U(\mathfrak{h})$ with p inverted. It follows from [2], Thm. 5.1.4 that the inclusion $U(\mathfrak{h}) \subseteq U(Lie(G_0^{p^m})) \subseteq D(G_0^{p^m})$ induces a canonical isomorphism of $\mathbb{Q}_p$-algebras

$$\widehat{U(\mathfrak{h})}_{\mathbb{Q}_p} \xrightarrow{\cong} D_{r_0}(G_0^{p^m}, \mathbb{Q}_p).$$

Indeed, by [loc.cit.] there is an explicit isomorphism between $\widehat{U(\mathfrak{h})}_{\mathbb{Q}_p}$ and the completion of $\mathbb{Q}_p\Lambda_{G_0^{p^m}} := \mathbb{Q}_p \otimes_{\mathbb{Z}_p} \mathbb{Z}_p[[G_0^{p^m}]]$ with respect to the norm on $\mathbb{Q}_p\Lambda_{G_0^{p^m}}$ constructed in [6], chap.7. By the explicit formulae given in [loc.cit.] this norm equals the restriction of $||.||^{(m)}_{r_0}$ to $\mathbb{Q}_p\Lambda_{G_0^{p^m}}$. But the latter space is norm-dense in $D_{r_0}(G_0^{p^m}, \mathbb{Q}_p)$ (cf. [19], p.19) and the resulting isomorphism is easily seen to be as claimed.

To ease notation we let

$$l := m + k$$

which is ≥ 1. We fix an ordered o_L-basis $\mathfrak{y}_1, ..., \mathfrak{y}_d$ of $\mathfrak{g}$ and recall our chosen $\mathbb{Z}_p$-basis $v_1, ..., v_n$ of o_L. The resulting $\mathbb{Z}_p$-basis of $\mathfrak{g}_0$ induces a positive PBW-filtration on the algebra $U(\mathfrak{g}_0)$ which makes it a *deformable* $\mathbb{Z}_p$*-algebra* in the language of [3], 3.6 with $U(\mathfrak{g}_0)_l := U(p^l\mathfrak{g}_0)$ being its *l-th deformation*. Denoting the p-adic completion of $U(\mathfrak{g}_0)_l$, with p inverted, by $\widehat{U(\mathfrak{g}_0)}_{l,\mathbb{Q}_p}$ we have, according to (3.7) a canonical isomorphism of $\mathbb{Q}_p$-algebras

$$\widehat{U(\mathfrak{h})}_{\mathbb{Q}_p} \xrightarrow{\cong} \widehat{U(\mathfrak{g}_0)}_{l,\mathbb{Q}_p}. \tag{3.8}$$

Alltogether we may write

$$D_{r_0}(G_0^{p^m}, L) = D_{r_0}(G_0^{p^m}, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} L = \widehat{U(\mathfrak{g}_0)}_{l,\mathbb{Q}_p} \otimes_{\mathbb{Q}_p} L =: \widehat{U(\mathfrak{g}_0)}_{l,L}. \tag{3.9}$$

Given $\alpha \in \mathbb{N}_0^{nd}$ we let $\mathfrak{Y}^\alpha := (v_1\mathfrak{y}_1)^{\alpha_{11}} \cdots (v_n\mathfrak{y}_d)^{\alpha_{nd}}$. Spelling out the definitions we observe that $\widehat{U(\mathfrak{g}_0)}_{l,L}$ equals the L-Banach algebra given by all series

$$\lambda := \sum_{\alpha \in \mathbb{N}_0^{nd}} d_\alpha \mathfrak{Y}^\alpha$$

where $d_\alpha \in L$ with $|d_\alpha| p^{l \cdot |\alpha|} \to 0$. It contains the kernel J of the map

$$\mathfrak{g}_0 \otimes_{\mathbb{Q}_p} L \to \mathfrak{g}, \ \mathfrak{x} \otimes a \mapsto a\mathfrak{x}.$$

If some algebra contains J we denote by $\langle J \rangle$ the two-sided ideal generated by J in this algebra. This should not cause any confusion. Reasoning as in (the proof of) [16], Prop. 2.4 the quotient algebra $\widehat{U(\mathfrak{g}_0)}_{l,L}/\langle J \rangle$ is given by the L-Banach algebra of all series

$$\lambda := \sum_{\beta \in \mathbb{N}_0^{d}} e_\beta \mathfrak{Y}'^\beta$$

where $e_\beta \in L$ with $|e_\beta| p^{l \cdot |\beta|} \to 0$ and $\mathfrak{Y}'^\beta := \mathfrak{y}_1^{\beta_1} \cdots \mathfrak{y}_d^{\beta_d}$. Reasoning as over $\mathbb{Z}_p$ this algebra coincides with $\widehat{U(\mathfrak{g})}_{el,L}$, i.e. with the π_L-adic completion, with π_L inverted, of the el-th deformation of the o_L-algebra $U(\mathfrak{g})$ endowed with its PBW-filtration relative to the basis $\mathfrak{y}_1, ..., \mathfrak{y}_d$. Recall that e equals the ramification index of L over $\mathbb{Q}_p$. Now taking into account the discussion preceding the map (2.3) we deduce a canonical isomorphism

$$D_{r_0}(G^{p^m}, L) = D_{r_0}(G_0^{p^m})/\langle J \rangle = \widehat{U(\mathfrak{g}_0)}_{l,L}/\langle J \rangle = \widehat{U(\mathfrak{g})}_{el,L}. \tag{3.10}$$

In particular, $D_{r_0}(G^{p^m}, K) = \widehat{U(\mathfrak{g})}_{el,L} \otimes_L K =: \widehat{U(\mathfrak{g})}_{el,K}$ as K-algebras. A this point we recall a deep theorem of [3].

Theorem 3.11. *(Ardakov/Wadsley) Let* $A := \widehat{U(\mathfrak{g})_{el,K}}$. *If* M *is a finitely generated (left)* A*-module with* $d_A(M) \geq 1$ *then* $d_A(M) \geq r$.

Proof. By choice of $\mathfrak{g}$ this follows from [loc.cit.], Thm. 9.9 combined with [loc.cit.], Lem. 2.6 since $el > 0$ and since K/L is finite. □

Remark: The theorem [loc.cit.], Thm. 9.9 is the analogue of Smith's theorem for p-adically completed deformations of universal enveloping algebras. It allows to prove a similar analogue for completed group algebras culminating in the main result of [loc.cit.].

By Cor. 3.4 we have

Corollary 3.12. *If* M *is a finitely generated (left)* $D_{r_m}(G)$*-module with* $d_{D_{r_m}(G)}(M) \geq 1$ *then* $d_{D_{r_m}(G)}(M) \geq r$.

3.13. We now deduce the locally analytic version of Smith's theorem. Let $\tilde{G}$ be an arbitrary locally L-analytic group of dimension d whose Lie algebra over L is split semisimple. Assume p is very good for this Lie algebra.

Let $\mathbf{G}$ be a connected split semisimple affine algebraic group scheme over o_L with Lie algebra $\mathfrak{g}$ such that $\mathfrak{g}_L := L \otimes_{o_L} \mathfrak{g}$ is isomorphic to $Lie(\tilde{G})$. Let us fix such an isomorphism. Recall that $\mathbf{G}$ gives rise to the natural number $r > 0$. Choose $k \geq 0$ sufficiently large and define a compact open subgroup G of $\tilde{G}$ which is L-uniform in the way described as above. The projective system of noetherian K-Banach algebras $D_{r_m}(G)$ for $m \geq 1$ defines the so-called *Fréchet-Stein structure* of $D(G)$. Without recalling the precise definition from [19] this essentially means that the natural map

$$D(G) \xrightarrow{\cong} \varprojlim_{m \geq 1} D_{r_m}(G)$$

is an isomorphism of topological K-algebras (where the right-hand side is endowed with the projective limit topology) and that the transition maps in the projective system are flat ring homomorphisms. This structure gives rise to a well-behaved abelian full subcategory $\mathcal{C}_{\tilde{G}}$ of the (left) $D(\tilde{G})$-modules, the *coadmissible modules*. By definition, an abstract $D(\tilde{G})$-module M is coadmissible if

(i) $M_m := D_{r_m}(G) \otimes_{D(G)} M$ is finitely generated over $D_{r_m}(G)$ for all $m \geq 1$;
(ii) the natural map $M \xrightarrow{\cong} \varprojlim_m M_m$ is an isomorphism.

The system $\{M_m\}_m$ is sometimes called the *coherent sheaf* associated to M. To give an example, any $D(\tilde{G})$-module which is finitely presented as $D(G)$-module is coadmissible.

Remark: The theory of Fréchet-Stein algebras is modelled according to the example $G = \mathbb{Z}_p$, the additive group of p-adic integers. In this case, the Fourier isomorphism of Y. Amice ([1]) identifies $D(\mathbb{Z}_p)$ with the ring of holomorphic functions on the rigid analytic open unit disc over K. The latter is a quasi-Stein space in the sense of R. Kiehl ([7]) and the Fourier isomorphism identifies the category of coadmissible modules with the coherent module sheaves on the disc.

By Prop. 3.3 the rings $D_{r_m}(G)$ are Auslander regular with global dimension equal to d. This implies ([19], §8) that the category $\mathcal{C}_{\tilde{G}}$ is equipped with a well-behaved *canonical codimension function* defined as $\mathrm{codim}(M) := j_{D(G)}(M)$ for a coadmissible module M. This number does not depend on the choice of G. To make the comparison with the work [3] more transparent we introduce the dimension function $\dim(M) := d - \mathrm{codim}\, M$. Hence $0 \leq \dim(M) \leq d$ if $M \neq 0$ and $\dim(0) = -\infty$.

For the definition of an *admissible locally analytic representation* as well as the foundations of the theory of such representations we refer to [19]. However, we at least recall ([loc.cit.], Thm. 6.3) that the abelian category $Rep^a_{\tilde{G}}(K)$ of admissible locally analytic $\tilde{G}$-representations over K is (essentially by definition of admissibility) anti-equivalent to $\mathcal{C}_{\tilde{G}}$. The functor is given by sending a representation V to its so-called *strong dual* $M := V'_b$. As a locally convex K-vector space V'_b equals the continuous dual of the locally convex vector space V endowed with a certain strong topology (see [17] for all terminology from non-archimedean functional analysis). The contragredient $\tilde{G}$-representation on V'_b extends naturally to a $D(\tilde{G})$-module structure giving the coadmissible module. It is known that any admissible-smooth $\tilde{G}$-representation is admissible ([loc.cit.], Thm. 6.6).
If V denotes an admissible locally analytic $\tilde{G}$-representation over K we let

$$\dim(V) := \dim(M)$$

where $M := V'_b$ is the associated coadmissible module.
It follows from [20], §5 that there is a filtration of the category $Rep^a_{\tilde{G}}(K)$ by Serre subcategories

$$Rep^a_{\tilde{G}}(K) = Rep^a_{\tilde{G}}(K)_d \supseteq Rep^a_{\tilde{G}}(K)_{d-1} \supseteq \ldots \supseteq Rep^a_{\tilde{G}}(K)_1 \supseteq Rep^a_{\tilde{G}}(K)_0$$

where $V \in Rep^a_{\tilde{G}}(K)_i$ if and only if $\dim(V) \leq i$. It is known ([15], Cor. 8.3) that the smooth-admissible representations are concentrated in dimension zero.

Theorem 3.14. *Let V denote an admissible locally analytic $\tilde{G}$-representation over K with* $\dim(V) \geq 1$. *Then* $\dim(V) \geq r$.

Proof. Let $M := V'_b$ denote the associated coadmissible module. As an immediate consequence of [19], lem. 8.4 we have $\dim(M) := \sup_{m \geq 1} d_{D_{r_m}(G)}(M_m)$ and hence $\dim(M) = d_{D_{r_{m'}}(G)}(M_{m'}) \geq 1$ for a particular $m' \geq 1$. Now apply cor. 3.12 to $M_{m'}$. □

Proposition 3.15. *Let V be as above. Then V is zero-dimensional if and only if the coherent sheaf of $M := V'_b$ consists of finite-dimensional K-vector spaces.*

Proof. If V is zero-dimensional then $d_{D_{r_m}(G)}(M_m) = 0$ for all $m \geq 1$ and the claim follows from prop. 3.3. The other implication is [15], thm. 8.4. □

REFERENCES

[1] Y. Amice. Duals. In *Proceedings of the Conference on p-adic Analysis (Nijmegen, 1978)*, volume 7806 of *Report*, pages 1–15. Katholieke Univ. Nijmegen, 1978.

[2] K Ardakov. Krull dimension of Iwasawa algebras and some related topics. *PhD-thesis. University of Cambridge (2004). Available at: http://www.maths.nottingham.ac.uk/personal/pmzka1/.*

[3] K. Ardakov and S. Wadsley. On irreducible representations of compact p-adic analytic groups. *preprint. Available at: http://arxiv.org/abs/1102.2606.*

[4] N. Bourbaki. *Éléments de mathématique. Fasc. XXXIII. Variétés différentielles et analytiques. Fascicule de résultats (Paragraphes 1 à 7).* Actualités Scientifiques et Industrielles, No. 1333. Hermann, Paris, 1967.

[5] David H. Collingwood and William M. McGovern. *Nilpotent orbits in semisimple Lie algebras.* Van Nostrand Reinhold Mathematics Series. New York, 1993.

[6] J. D. Dixon, M. P. F. du Sautoy, A. Mann, and D. Segal. *Analytic pro-p groups*, volume 61 of *Cambridge Studies in Advanced Mathematics.* Cambridge University Press, Cambridge, second edition, 1999.

[7] R. Kiehl. Theorem A und Theorem B in der nichtarchimedischen Funktionentheorie. *Invent. Math.*, 2:256–273, 1967.

[8] J. Kohlhaase. Invariant distributions on p-adic analytic groups. *Duke Math. J.*, 137(1):19–62, 2007.

[9] M. Lazard. Groupes analytiques p-adiques. *Inst. Hautes Études Sci. Publ. Math.*, (26):4–219, 1965.

[10] Thierry Levasseur. Some properties of noncommutative regular graded rings. *Glasgow Math. J.*, 34(3):277–300, 1992.

[11] H. Li and F. van Oystaeyen. *Zariskian filtrations*, volume 2 of *K-Monographs in Mathematics.* Kluwer Academic Publishers, Dordrecht, 1996.

[12] J. C. McConnell and J. C. Robson. *Noncommutative Noetherian rings.* Pure and Applied Mathematics (New York). John Wiley & Sons Ltd., Chichester, 1987.

[13] S. Orlik and M. Strauch. On the irreducibility of locally analytic principal series representations. *Represent. Theory*, 14:713–746, 2010.

[14] D.S. Passman. *Infinite crossed products*, volume 135 of *Pure and Applied Mathematics.* Academic Press Inc., Boston, MA, 1989.

[15] T. Schmidt. Auslander regularity of p-adic distribution algebras. *Represent. Theory*, 12:37–57, 2008.

[16] T. Schmidt. Analytic vectors in continuous p-adic representations. *Compositio Math.*, 145:247–270, 2009.

[17] P. Schneider. *Nonarchimedean functional analysis.* Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2002.

[18] P. Schneider. Continuous representation theory of p-adic Lie groups. In *International Congress of Mathematicians. Vol. II*, pages 1261–1282. Eur. Math. Soc., Zürich, 2006.

[19] P. Schneider and J. Teitelbaum. Algebras of p-adic distributions and admissible representations. *Invent. Math.*, 153(1):145–196, 2003.

[20] P. Schneider and J. Teitelbaum. Duality for admissible locally analytic representations. *Represent. Theory*, 9:297–326 (electronic), 2005.

[21] S.P. Smith. Krull dimension of factor rings of the enveloping algebra of a semisimple Lie algebra. *Math. Proc. Cambridge Philos. Soc.*, 93(3):459–466, 1983.

Mathematisches Institut, Westfälische Wilhelms-Universität Münster, Einsteinstr. 62, D-48149 Münster, Germany
E-mail address: `toschmid@math.uni-muenster.de`